

Screening for possible presence of a coexisting mental health disorder in members who have a substance use disorder

SCREENING NAME	WHO IS ELIGIBLE FOR SCREENING	CONDITIONS WHEN IT SHOULD BE APPLIED	AGE RANGE	# OF QUESTIONS	WHO COMPLETES	FREQUENCY OF ADMINISTRATION	WHERE TO ACCESS
CHILD BEHAVIOR CHECKLIST (CBCL/6-18)	Children and adolescents (6-18 year olds)	To assess grief/loss and anxiety/mood	6-18 year olds	120	Parents or caregiver	Every 6 months	Child Behavior Checklist for Ages 6-18
PARENT STRESS SCALE	Parents of children	Measures level of stress by parents- both positive and negative aspects of parenting	Adult	18	Parents or caregiver	Pre and Post Treatment	Parenting Stress Scale
SNAP-IV	Children and adolescents (6-18) year olds	To screen children and adolescents for ADHD	6-18 year olds	18	Teacher and parent	No recommendation provided	SNAP-IV Teacher and Parent Rating Scale
PHQ-9	Available in English, Spanish and many other languages	Any suspicion of depression; good practice for primary care screenings	18+ but has been used in children as young as 12	9	Patient	No limit	Patient Health Questionnaire-9
GAD-7	Available in English and Spanish	Any suspicion that anxiety is present or playing a role in presenting problem	18+; but has been used in 11-17 year olds	7	Patient	No limit	Generalized Anxiety Disorder 7-item (GAD-7) scale
MDQ	Outpatient setting; Primary care patients receiving treatment for depression	To screen adults for bipolar disorder	Adult	15	Patient	No recommendation provided	Mood Questionnaire
PC-PTSD-5	Any Adult	Designed for universal use in VA primary care settings	Tested in Adults	5	Clinician or patient	No recommendation provided	PC-PTSD-5
MHSF-III	Adults in SUD settings	Designed for MH screening in SUD settings	Tested in Adults	18 'yes/no' questions	Clinician or patient	Screening upon admission to SUD programs	Mental Health Screening Form-III (MHSF-III)

SCREENING NAME	WHO IS ELIGIBLE FOR SCREENING	CONDITIONS WHEN IT SHOULD BE APPLIED	AGE RANGE	# OF QUESTIONS	WHO COMPLETES	FREQUENCY OF ADMINISTRATION	WHERE TO ACCESS
KESSLER-6 DISTRESS SCALE	Any adult; available in multiple languages	To distinguish SMI (serious mental illness) cases from non-SMI cases	Validated for use with adults	6	Clinician or patient (there are self-administered and interviewer-administered versions)	No recommendation provided	K-6 Scales
KESSLER- 10 DISTRESS SCALE	Any adult; available in multiple languages	To distinguish SMI (serious mental illness) cases from non-SMI cases	Validated for use with adults	10	Clinician or patient (there are self-administered and interviewer-administered versions)	No recommendation provided	K-10 Scales

